

## ARTICLE

# Examination of Surface Water Along River-Rima Floodplain in Wamakko, Sokoto State, Nigeria

Abdulqadir Abubakar Usman<sup>1</sup> Murtala Abubakar Gada<sup>2</sup> Aminu Muhammad Bayawa<sup>3</sup>  
Ibrahim Mustapha Dankani<sup>2</sup> Saadu Umar Wali<sup>1\*</sup>

1. Department of Geography, Federal University Birnin-kebbi, Kebbi State, Nigeria

2. Department of Geography, Usmanu Danfodiyo University Sokoto, Sokoto State, Nigeria

3. Department of Pure and Applied Chemistry, Usmanu Danfodiyo University Sokoto, Sokoto State, Nigeria

### ARTICLE INFO

#### Article history

Received: 14 July 2020

Accepted: 29 July 2020

Published Online: 30 July 2020

#### Keywords:

Surface water pollution

Agricultural activities

Bacteriological analysis.

### ABSTRACT

This study examined the hydrochemistry of surface water along the River-Rima floodplain area. Five sampling locations were purposively selected, and, in each point, three samples were taken from surface water (river). The sampling was repeated after 20 days. Thus, a total of 30 samples were collected. Water samples obtained were subjected to laboratory tests. Results revealed that BOD, TDS,  $Mg^{2+}$ , and  $Fe^{3+}$  are above the World Health Organization (WHO) and Standard Organization of Nigeria (SON) reference guidelines for drinking water quality. Isolates detected from the coliform bacteriological analysis include *Enterobacter aerogene*, *Escherichia coli*, and *Citrobacter freundii* with most of the samples showing coliform bacteria growth above the SON standard for drinking water. Hence, the water in the River-Rima floodplain of the Wamakko area is of low quality and unsafe for drinking. Results of principal component analysis (PCA) revealed external influences such as pollutant wash off and rock weathering as controls on hydrochemistry of surface water. There is some indication of anthropogenic inputs ( $Cl^-$ ,  $NO_3^-$ , and  $PO_4^{2-}$ ) based on hierarchical cluster analysis. Elements including  $Cl^-$ ,  $NO_3^-$ , and  $PO_4^{2-}$  are increasingly added into surface water from human activities, mainly agriculture, and municipal sewage.

## 1. Introduction

Freshwater resource deterioration is now a global problem and is increasing at a faster rate<sup>[1-3]</sup>. According to<sup>[4]</sup> one of the most critical crises in developing countries is the lack of adequate potable water. In some remote communities, surface water is consumed raw with little or no water treatment. Surface water is easily polluted and requires treatment before use<sup>[5-9]</sup>.

Water pollution occurs when water becomes overloaded with chemicals and the aquatic organisms cannot keep up with their cleaning responsibilities<sup>[10-12]</sup>. Some may die and others may grow too fast. There are many types and sources of pollutants that can pollute surface and groundwater<sup>[13-17]</sup>. The use of fertilizers and pesticides on cropland, gardens, and yards help farms and homeowners, but can also be damaging to aquatic and marine life when not properly applied or managed<sup>[18-20]</sup>. Poorly managed

\*Corresponding Author:

Saadu Umar Wali,

Department of Geography, Federal University Birnin kebbi, P.M.B 1157. Kebbi State, Nigeria;

Email: [saadu.umar@fubk.edu.ng](mailto:saadu.umar@fubk.edu.ng)



gold, kaolin, gypsum, and marble.

### 3. Materials and Methods

#### 3.1 Sample Collection

This study covers Kwalkwalawa, Marmaro, Asare, Boye, and Kaura Abdu villages. Five sampling points were selected (Figure 1). Global Positioning System (GPS) was used to locate the sampling points. At each point, three samples were taken from the surface water. The sampling was repeated after 20 days. Thus, a total of 30 samples were collected for laboratory analysis. The sampling was carried out between October and November 2015 at the cessation of rains, to avoid the influence of runoff. At the sampling points, the sample bottles were rinsed with the sample water twice before water samples were taken. The bottles were corked tightly and labeled for identification and taken to the laboratory.

#### 3.2 Laboratory Analysis

The Physicochemical analyses conducted include the determination of pH (Digital pH meter), Dissolved Oxygen (DO) (Iodometric titration method), Biochemical Oxygen Demand (BOD) (Incubation bottles), Total Dissolved Solids (TDS) (Glass fiber filter method), Total Suspended Solids (TSS) (Filter paper method), Chloride (Cl<sup>-</sup>) (Titration method), Phosphate (PO<sub>4</sub><sup>3-</sup>) (Spectrophotometer), Nitrate (NO<sub>3</sub><sup>-</sup>) (Macro – Kjeldahl flask method), Potassium (K<sup>+</sup>) (Flame photometer), Magnesium (Mg<sup>2+</sup>) (Titration method), Calcium (Ca<sup>2+</sup>) (Titration method) and Iron (Fe<sup>3+</sup>) (Atomic Absorption Spectrometer). The WHO and SON standards were used to compare the levels of pollutants for drinking purposes. The bacteria indicators of surface water pollution were analyzed using the Most Probable Number (MPN) to find the number of coliforms. Isolate method was also used to identify and detect various coliform bacteria found in the water samples.

#### 3.3 Statistical Analysis

Data obtained from the laboratory tests were summarized (mean±standard deviation) and standardized. This was followed by multivariate analysis (Principal Component Analysis; and Hierarchical Cluster Analysis). PCA was used to reduce the studied parameters to more interpretable form<sup>[60-64]</sup>. It is used to reduce huge and complex datasets by finding a small number of variables that define the utmost variance. PCA was carried out on a subset of 11 chosen hydrochemical parameters (pH, DO, BOD, TDS, TSS, Cl<sup>-</sup>, PO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>3+</sup>), which characterized the general groundwater composition. Hierarchical

cluster analysis (HCA), was conducted to classify sampling locations<sup>[65-67]</sup>. In clustering, locations with comparable surface water composition or otherwise parameters with typical properties would be gathered into identical clusters<sup>[60-64]</sup>. Ward's-algorithmic grouping technique after the Euclidean distance was used. This method does not account for data validity and disconnected values and neglects credible and possible connections or resemblances (statistical meddling) between variables, which are somewhat clear in most hydrochemical data<sup>[60-64]</sup>.

### 4. Results and Discussion

#### 4.1 Physicochemical Characteristics

Tables 1 and 2 summarized the results (mean±standard deviation) physicochemical and bacteriological analysis. The mean values for the sampling locations were compared with the World Health Organization (WHO), and the Standards Organization of Nigeria (SON) standards. The pH level ranged between 6.6 -7.1. The highest mean concentration of 7.1± 0.1 was detected at Kaura Abdu and the lowest is 6.6 ± 0.1 was detected at Kwalkwalawa. Since pH is one of the most significant operational water quality parameters, (its range affects the workability of other parameters), moderate pH levels are required (Duvbiana and Egbuna, 2013). The pH values from all the five (5) locations are within the WHO reference limits of 6.5- 8.5.

Dissolved Oxygen (DO) ranged from 5.7 to 7.4 mg/l. The highest concentration of DO was observed in Asare and the lowest level was observed in Marmaro. The DO concentration was generally within the WHO reference limits, indicating water of good quality. Low DO is evidence of reduced bacterial activity within the water ecosystem. Biological Oxygen Demand (BOD) ranged from 23.4 mg/l to 14.7 mg/l. The BOD levels for all sites were above the WHO maximum permissible limits, indicating water that is unsafe for drinking. The low value for BOD at Boye might be due to low organic matter, while that for Kaura Abdu water might be due to a high concentration of pollutants and organic matter. The mean concentrations of BOD values were higher compared to the values recorded by<sup>[68]</sup> and<sup>[69]</sup> who recorded lower values (7.38 mg/l and 0.7 mg/l respectively).

Total Dissolved Solids (TDS) ranged from 7.2 mg/l to 2.2. The mean concentrations indicated that all the sampled locations had TDS values below the SON maximum allowable limits (500 mg/l). Dissolved solids in water can be used water as an indication of its suitability for domestic use<sup>[70]</sup>. TDS in drinking water is not of health

**Table 1.** Summary of Physicochemical parameters

Sampling	Parameters											
	Points	pH	DO	BOD	TDS	TSS	Cl <sup>-</sup>	PO <sub>4</sub> <sup>3-</sup>	NO <sub>3</sub> <sup>-</sup>	K <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>
Kwawkwalawa	6.6 ± 0.1	7.1 ± 1.1	20.1 ± 1.4	7.2 ± 2.5	5.8 ± 0.3	0.8 ± 0.2	0.2 ± 0.0	1.1 ± 0.1	1.9 ± 0.1	0.7 ± 0.1	0.4 ± 0.0	4.5 ± 0.0
Marmaro	6.7 ± 0.1	5.7 ± 2.0	16.7 ± 3.6	5.3 ± 0.6	3.3 ± 0.3	1.1 ± 0.3	0.2 ± 0.0	1.0 ± 0.1	2.0 ± 0.1	0.7 ± 0.0	0.5 ± 0.1	2.0 ± 0.0
Share	6.8 ± 0.0	7.4 ± 2.3	16.8 ± 3.9	5.2 ± 0.8	5.2 ± 0.3	1.1 ± 0.1	0.2 ± 0.0	1.2 ± 0.2	2.1 ± 0.1	0.8 ± 0.1	0.5 ± 0.1	2.0 ± 0.0
Boye	6.7 ± 0.0	7.1 ± 1.3	14.7 ± 1.3	3.2 ± 0.6	2.3 ± 0.6	1.4 ± 0.3	0.2 ± 0.0	1.6 ± 0.2	2.0 ± 0.0	0.7 ± 0.0	0.4 ± 0.0	2.1 ± 0.0
Kaura Abdu	7.1 ± 0.1	7.1 ± 0.6	23.4 ± 1.7	2.2 ± 0.3	1.2 ± 0.3	1.1 ± 0.1	0.2 ± 0.0	0.9 ± 0.1	2.0 ± 0.1	1.2 ± 0.1	0.6 ± 0.0	0.3 ± 0.0
WHO	6.5-8.5	7.5	10	1	10	250	5	10	250	250	-	0.3
SON	6.5-8.5	-	-	500	-	250	-	50	-	-	0.2	0.3

concern, but it affects palatability levels above 600 mg/l<sup>[71]</sup>. Total Suspended Solids (TSS) were high at Kwawkwalawa (5.8 ± 0.3 mg/l) perhaps due to high silt and organic matter contents. Low values were recorded at Kaura Abdu (1.2±0.3). Chloride (Cl<sup>-</sup>). The highest concentration was observed at Boye (1.4±0.3 mg/l). Chloride increases with the increasing degree of eutrophication<sup>[72,73]</sup>. The lowest values recorded in this study are from Kwawkwalawa (0.8±0.2 mg/l). All the samples showed satisfactory levels of WHO and SON standard limits.

The phosphate ion (PO<sub>4</sub><sup>3-</sup>) concentration was low. Values obtained were within the maximum allowable limits by the WHO (5 mg/l). The highest concentrations of 0.2±0.0 mg/l were observed for all the surface samples. Phosphate occurs naturally in low concentrations in natural water while higher concentrations may pollution from human activities<sup>[70]</sup>. Nitrate (NO<sub>3</sub><sup>-</sup>) ranged from 0.9 mg/l to 1.6 mg/l. The concentrations were below the WHO and SON permissible limits. Nitrate is mainly derived from organic matter and oxidation of ammonia. However, NO<sub>3</sub><sup>-</sup> is increasingly added to surface water from chemical fertilizers. The effects of leached fertilizers on water environments and their health implications have been reported<sup>[74,75]</sup>.

Potassium Ion (K<sup>+</sup>) ranged from 2.1 mg/l to 1.9 mg/l. Both samples have potassium concentrations below the WHO maximum permissible limits of 250 mg/l. Calcium Ion (Ca<sup>2+</sup>) ranged from 1.2 mg/l 0.7 mg/l. Ca<sup>2+</sup> concentration is below the maximum allowable WHO limits (250 mg/l). Calcium is an indication of hardness in water, but it has no known effect on the health of human beings. Magnesium (Mg<sup>2+</sup>) ranged from 0.6 to 0.4 mg/l. The concentrations were above the SON limit of 0.2mg/l. A high concentration of Mg<sup>2+</sup> gives an unpleasant taste to drinking water<sup>[75]</sup>. Iron (Fe<sup>3+</sup>) was detected in appreciable quantities. The highest concentration was found in Kwawkwalawa (4.5 ± 0.0), and the lowest was found in Kaura Abdu (0.3). These samples are above the WHO and SON allowable limits.

## 4.2 Bacteriological Analysis

Coliform growth was higher in all samples. The three coliform bacteria detected are *Citrobacter freundii*, *Enterobacter aerogene*, and *Escherichia coli*. Table 2 shows the rate of contamination of surface water concerning coliform bacteria isolated from surface water samples. The rate of contamination of surface water samples as presented in Table 2 showed Kwawkwalawa had 655, Marmaro 625, Asare 625, Boye 180, and Kaura Abdu 26, respectively. In this study, all five samples showed a positive presumptive test. Therefore, the entire samples collected were not suitable for drinking because concentrations exceed the rate of 10 as recommended by<sup>[76]</sup>. Water samples collected from Kwawkwalawa have the highest rate of contamination compared to the other sample. Therefore, a potential health risk exists due to the presence of microbial pathogens in the water. The *Enterobacter aerogene* population was 33.33%, *Escherichia coli* had 11.11%, while *Citrobacter freundii* had 55.6%, respectively. The main sources of these microbial contaminants in wastewater are human and animal wastes.

**Table 2.** Mean MPN of Coliforms and Bacteria Isolated from surface water

Kwawkwalawa	655	<i>Enterobacter aerogene</i> & <i>Citrobacter freundii</i>
Marmaro	625	<i>Citrobacter freundii</i> & <i>Enterobacter aerogene</i>
Share	625	<i>Citrobacter freundii</i>
Boye	180	<i>Enterobacter aerogene</i> & <i>Escherichia coli</i>
Kaura Abdu	26	<i>Citrobacter freundii</i>

## 4.3 Multivariate Analysis


### 4.3.1 Principal Component Analysis

Principal component analysis (PCA) is a valuable statistical tool that can be used to reduce hydrochemical data and relates it to sources of solutes in surface water. PCA was carried out on 13 subsets of elements (Table 3). Com-

ponent 1 explained 58.63 % of the total variance and had high positive loadings ( $\geq 0.65$ ) on BOD. Component 2 had moderate loading on BOD, TDS, and TSS, it explained 36.08 % of the total variance. Significant loading on DO occurred in Component 3 with moderate loading on TSS. Components 1-3 relate to external influences on surface water quality. Component 4 had significant loading on  $Fe^{3+}$ . This component can be related to rock weathering. High  $Fe^{3+}$  concentrations exceeding WHO reference guidelines in surface and groundwater were reported from different parts of the Sokoto basin [77-80]. Figure 2 further illustrates a biplot of PC 1 and PC 2. It is difficult to identify any pattern.

**Table 3.** Principal components (PCs) of hydrochemical data

Variables	PC 1	PC 2	PC 3	PC 4
pH	0.046	-0.025	0.055	-0.077
DO	0.030	0.031	0.759	0.283
BOD	0.873	0.476	-0.019	0.022
TDS	-0.291	0.570	-0.376	-0.204
TSS	-0.300	0.537	0.490	-0.394
$Cl^-$	-0.011	-0.072	0.035	0.035
$PO_4^{2-}$	0.000	0.000	0.000	0.000
$NO_3^-$	-0.048	-0.042	0.117	0.198
$K^+$	-0.003	-0.013	0.042	-0.070
$Ca^{2+}$	0.057	-0.014	0.055	-0.049
$Mg^{2+}$	0.018	-0.007	-0.001	-0.073
$Fe^{3+}$	-0.231	0.388	-0.138	0.814
% variance	58.626	36.075	3.061	2.238
Eigenvalue	13.033	8.020	0.680	0.498




**Figure 2.** Further illustrates a biplot of PC 1 and PC 2

### 4.3.2 Hierarchical Cluster Analysis

Hierarchical cluster analysis (HCA) is a commonly

used procedure in water quality analysis since it is capable of differentiating sampling locations [65-67,81-86]. By means, HCA, the sampling locations with comparable characteristics can be grouped into a cluster. The graphic assembly of the grouping process is shown in Figure 3. Cluster 1 is comprised of sampling locations with similar concentrations of pH,  $Ca^{2+}$ ,  $Mg^{2+}$ , and BOD. Cluster 2 is comprised of locations with similar concentrations of DO,  $K^+$ ,  $Cl^-$ ,  $PO_4^{3-}$  and  $NO_3^-$ . Cluster 3 contained locations with similar TDS, TSS, and  $Fe^{3+}$  concentrations. Cluster 2 can be related to anthropological influences, since  $Cl^-$ ,  $PO_4^{2-}$  and  $NO_3^-$  are increasingly added into surface water by human activities [87,88].



**Figure 3.** Dendrogram produced from cluster analysis based on the sampling locations to identify the major surface water characteristics

## 5. Conclusion

This study aimed at examining the hydrochemistry of surface water along River-Rima floodplain at Wamakko, Sokoto State, Nigeria. The study discovered that BOD, TDS,  $Mg^{2+}$ , and  $Fe^{3+}$  are not following the WHO and SON standards. The presence of  $Cl^-$ ,  $NO_3^-$ , and  $PO_4^{3-}$  in detectable quantities, indicated high toxicity levels in surface water. Coliform and total bacteria count indicated microbial pollution from anthropogenic activities. The isolates bacteria detected include *Enterobacter aerogene*, *Escherichia coli*, and *Citrobacter freundii*. The presence of *E. coli* in detectable quantities indicated high contamination and therefore posed a serious risk to human and animal health. Finally, the quality of the water is deteriorating due to the continuous contamination of the water by farmers in the process of fertilizer application, dumping of agricultural waste, and defecation by animals. Hence, the water in the Sokoto River Rima floodplain of the Wamakko local area of Sokoto State is of low standard and unsafe for drinking.

## Acknowledgments

This study was supported by Federal University Birnin Kebbi. Thanks to all anonymous contributors.

## References

- [1] Mahananda, H.B., M.R. Mahananda, B.P. Mohanty. Studies of Physico-Chemical and Biological parameters of a Freshwater Pond Ecosystem as an Indicator of Water Pollution. *Ecological Environmental and Conservation*, 2005, 11(3-4): 537-541.
- [2] Aboyeji, O.O. Freshwater pollution in some Nigerian local communities, causes, consequences and probable solutions. *Academic Journal of Interdisciplinary Studies*, 2013, 2(13): 111-117.
- [3] Chen, J., et al. Challenges and prospects of sustainable groundwater management in an agricultural plain along the Silk Road Economic Belt, north-west China. *International Journal of Water Resources Development*, 2016, 34(3): 354-368.
- [4] Efe, S.I., et al. Seasonal variations of physico-chemical characteristics in water resources quality in Western Niger Delta Region, Nigeria. *Journal of Applied Sciences and Environmental Management*, 2005, 9(1): 191 - 195.
- [5] Hunter, P.R., D. Zmirou-Navier, P. Hartemann. Estimating the impact on health of poor reliability of drinking water interventions in developing countries. *Sci Total Environ*, 2009, 407(8): 2621-4.
- [6] Lea, M. Bioremediation of Turbid Surface Water Using Seed Extract from the *Moringa oleifera* Lam. (Drumstick) Tree. *Curr Protoc Microbiol*, 2014, 33: 1G 2 1-8.
- [7] Mwabi, J.K., et al. Household water treatment systems: A solution to the production of safe drinking water by the low-income communities of Southern Africa. *Physics and Chemistry of the Earth, Parts A/B/C*, 2011, 36(14-15): 1120-1128.
- [8] Mwabi, J.K., B.B. Mamba, M.N. Momba. Removal of *Escherichia coli* and faecal coliforms from surface water and groundwater by household water treatment devices/systems: a sustainable solution for improving water quality in rural communities of the Southern African development community region. *Int J Environ Res Public Health*, 2012, 9(1): 139-70.
- [9] Naidoo, S., A.O. Olaniran. Treated wastewater effluent as a source of microbial pollution of surface water resources. *Int J Environ Res Public Health*, 2013, 11(1): 249-70.
- [10] Azizullah, A., et al. Water pollution in Pakistan and its impact on public health--a review. *Environ Int*, 2011, 37(2): 479-97.
- [11] Tang, J.Y., et al. Toxicity characterization of urban stormwater with bioanalytical tools. *Water Res*, 2013, 47(15): 5594-606.
- [12] Yuan, Z., et al. Sweating the assets - The role of instrumentation, control and automation in urban water systems. *Water Res*, 2019, 155: 381-402.
- [13] Dinka, M.O., W. Loiskandl, J.M. Ndambuki. Hydrochemical characterization of various surface water and groundwater resources available in Matahara areas, Fantalle Woreda of Oromiya region. *Journal of Hydrology: Regional Studies*, 2015, 3: 444-456.
- [14] Andrade, L., et al. Surface water flooding, groundwater contamination, and enteric disease in developed countries: A scoping review of connections and consequences. *Environ Pollut*, 2018, 236: 540-549.
- [15] Li, J., F. Li, Q. Liu. PAHs behavior in surface water and groundwater of the Yellow River estuary: Evidence from isotopes and hydrochemistry. *Chemosphere*, 2017, 178: 143-153.
- [16] Pavlidis, G., V.A. Tsihrintzis. Environmental Benefits and Control of Pollution to Surface Water and Groundwater by Agroforestry Systems: a Review. *Water Resources Management*, 2017, 32(1): 1-29.
- [17] Singh, J., et al. *Water Pollutants: Origin and Status*. 2020: 5-20.
- [18] Gorde, S.P., M.V. Jadhav. Assessment of water quality parameters: A review. *Journal of Engineering Research and Applications*, 2013, 3(6): 2029-2035.
- [19] Chartzoulakis, K., M. Bertaki. Sustainable Water Management in Agriculture under Climate Change. *Agriculture and Agricultural Science Procedia*, 2015, 4: 88-98.
- [20] Kaur, T., R. Bhardwaj, S. Arora. Assessment of groundwater quality for drinking and irrigation purposes using hydrochemical studies in Malwa region, southwestern part of Punjab, India. *Applied Water Science*, 2016, 7(6): 3301-3316.
- [21] Rim-Rukeh, A., O.G. Ikhifa, A.P. Okokoyo. Effects of Agricultural Activities on the Water Quality of Orogo River, Agbor Nigeria. *Journal of Applied Sciences Research*, 2006, 2(5): 256-259.
- [22] Clemens, M., et al.. Groundwater protection under water scarcity; from regional risk assessment to local wastewater treatment solutions in Jordan. *Sci Total Environ*, 2020, 706: 136066.
- [23] Gwenzi, W., N. Chaukura. Organic contaminants in African aquatic systems: Current knowledge, health risks, and future research directions. *Sci Total Environ*, 2018, 619-620: 1493-1514.
- [24] Kumar, A., D. Pal. Antibiotic resistance and wastewater: Correlation, impact and critical human health challenges. *Journal of Environmental Chemical En-*

- gineering, 2018, 6(1): 52-58.
- [25] Moeder, M., et al. Potential of vegetated ditches to manage organic pollutants derived from agricultural runoff and domestic sewage: A case study in Sinaloa (Mexico). *Sci Total Environ*, 2017, 598: 1106-1115.
- [26] Gaballah, M.S., Khalaf, K., Beck, A., Lopez, J. Water Pollution concerning Agricultural Activity Impact in Egypt. *Journal of Applied Science Research*, 2005, 1: 9-17.
- [27] Naubi, I., et al. Effectiveness of Water Quality Index for Monitoring Malaysian River Water Quality. *Polish Journal of Environmental Studies*, 2016, 25(1): 231-239.
- [28] Lkr, A., M.R. Singh, N. Puro. Assessment of water quality status of Doyang River, Nagaland, India, using Water Quality Index. *Applied Water Science*, 2020, 10(46): 1-13.
- [29] Boukari, A., et al. Assessment of Agriculture Pressures Impact on the Joumine River Water Quality Using the PEGASE Model. *Environ Manage*, 2019, 64(4): 520-535.
- [30] Clément, F., et al. Landscape diversity and forest edge density regulate stream water quality in agricultural catchments. *Ecological Indicators*, 2017, 72: 627-639.
- [31] Conroy, E., et al. The impact of cattle access on ecological water quality in streams: Examples from agricultural catchments within Ireland. *Sci Total Environ*, 2016, 547: 17-29.
- [32] Diamantini, E., et al. Driver detection of water quality trends in three large European river basins. *Sci Total Environ*, 2018, 612: 49-62.
- [33] Evans, A.E.V., et al. Agricultural water pollution: key knowledge gaps and research needs. *Current Opinion in Environmental Sustainability*, 2019, 36: 20-27.
- [34] Razali, A., et al. Land use change in highland area and its impact on river water quality: a review of case studies in Malaysia. *Ecological Processes*, 2018, 7(19): 1-17.
- [35] Taylor, S.D., Y. He, K.M. Hiscock. Modelling the impacts of agricultural management practices on river water quality in Eastern England. *J Environ Manage*, 2016, 180: 147-63.
- [36] Van der Laan, M., R. Van Antwerpen, K.L. Bristow. River water quality in the northern sugarcane-producing regions of South Africa and implications for irrigation: A scoping study. *Water SA*, 2012, 38(1): 87-96.
- [37] EPA. Environmental Protection Agency: Distribution system indicators of drinking water quality, 2006: 1-100.  
<https://pdfs.semanticscholar.org/e989/3a212f9e-ae94dffce9dc10cb4fbc1c1bd02d.pdf>
- [38] Barmntlo, S.H., et al. Assessing combined impacts of agrochemicals: Aquatic macroinvertebrate population responses in outdoor mesocosms. *Sci Total Environ*, 2018, 631-632: 341-347.
- [39] Horn, S., R. Pieters, T. Bohn. May agricultural water sources containing mixtures of agrochemicals cause hormonal disturbances? *Sci Total Environ*, 2020, 711: 134862.
- [40] Maishanu, H.M., et al. Influence of Human Activities on the Phytoplankton Communities of River Rima in Kwalkwalwa Area of Sokoto State, Nigeria. *Science World Journal*, 2019, 14(3): 42-47.
- [41] Adejuwon, J.O. Assessment of the Changing Pattern in Maize Cultivation in Sokoto-Rima River Basin, Nigeria. *Journal of Applied Sciences and Environmental Management*, 2018, 22(9): 1433.
- [42] Emeribe, C.N., E.T. Ogbomida, J.O. Enoma-Calus. Climatic Variability and Estimation of Supplementary Irrigation Water Needs of Selected Food Crops in the Sokoto-Rima River Basin, Nigeria. *Nigerian Journal of Environmental Sciences and Technology*, 2019, 3(1): 86-104.
- [43] Eniolorunda, N.B., S.A. Mashi, G.N. Nsofor. Toward achieving a sustainable management: characterization of land use/land cover in Sokoto Rima floodplain, Nigeria. *Environment, Development and Sustainability*, 2016, 19(5): 1855-1878.
- [44] Simeonov, V., et al. Assessment of the surface water quality in Northern Greece. *Water Research*, 2003, 37(17): 4119-4124.
- [45] Giri, S., A.K. Singh. Risk assessment, statistical source identification and seasonal fluctuation of dissolved metals in the Subarnarekha River, India. *J Hazard Mater*, 2014, 265: 305-14.
- [46] Akkoyunlu, A., M.E. Akiner. Pollution evaluation in streams using water quality indices: A case study from Turkey's Sapanca Lake Basin. *Ecological Indicators*, 2012, 18: 501-511.
- [47] Varol, M., B. Şen. Assessment of nutrient and heavy metal contamination in surface water and sediments of the upper Tigris River, Turkey. *Catena*, 2012, 92: 1-10.
- [48] Abdullahi, S.A., et al. Assessment of water availability in the Sokoto Rima River Basin. *Resources and Environment*, 2014, 4(4): 220-233.
- [49] Ita, E.O. Inland Fishery Resources of Nigeria. CIFA Occasional Paper No. 2 EAO, Rome Italy. 1993: 1-120.
- [50] NIMET. Climate review: Nigerian Meteorological Agency. 2012.
- [51] Adegboyega, S.A., et al. Monitoring drought and ef-

- fects on vegetation in Sokoto state, Nigeria using statistical and geospatial techniques. *Ethiopian Journal of Environmental Studies and Management*, 2016, 9(1): 56-69.
- [52] Ekoh, H.C. Analysis of rainfall trend in Sokoto State, Nigeria (1987-2016). *World News of Natural Sciences*, 2020, 28: 171-186.
- [53] Wali, S.U., et al. Hydrochemical characterization of shallow and deep groundwater in Basement Complex areas of southern Kebbi State, Sokoto Basin, Nigeria. *Applied Water Science*, 2019, 9(169): 1-36.
- [54] Hayatu, N.G., et al. Effect of land utilization types on some physical and hydraulic properties of soils in Gidan Sule, Wamakko, Sokoto State, Nigeria. *International Journal of Research and Innovation in Applied Science*, 2018, 3(11): 14-18.
- [55] Davis, G. Rainfall and temperature: Sokoto State in maps. *An Atlas Physical and Human Resources*. John Wiley and Sons. Inc., New York. 1984: ATLAS MAP.
- [56] Akinbiyi, O.A., et al. Geophysical characterization and hydraulic properties of unconsolidated floodplain aquifer system in Wamako area, Sokoto State, north-western Nigeria. *Applied Water Science*, 2019, 9(177): 1-10.
- [57] Wali, U. Surface and Groundwater: The Impact of Climate Change on Sokoto State: Evidence and Challenges. UNDP/Sokoto State Government. Eds: M.A. Aliya and A.G. Fada. 2013: 1-151.
- [58] Hamidu, H., et al. Groundwater Resource Appraisals of Bodinga and Environs, Sokoto Basin North Western Nigeria. *Nigerian Journal of Basic and Applied Sciences*, 2017, 24(2): 92.
- [59] Mamman, A.B. Nigeria; Giant in the Tropics. Gabumo Publishing Company, 2nd Edition, Abuja, Nigeria. 2000: 1-100.
- [60] Lin, C.Y., et al. Delineation of temporal variability and governing factors influencing the spatial variability of shallow groundwater chemistry in a tropical sedimentary island. *Journal of Hydrology*, 2012, 432-433: 26-42.
- [61] Azhar, S.C., et al. Classification of river water quality using multivariate analysis. *Procedia Environmental Sciences*, 2015, 30: 79-84.
- [62] Blake, S., et al. Compositional multivariate statistical analysis of thermal groundwater provenance: A hydrogeochemical case study from Ireland. *Applied Geochemistry*, 2016, 75: 171-188.
- [63] Rakotondrabe, F., et al. Water quality assessment in the Betare-Oya gold mining area (East-Cameroon): Multivariate Statistical Analysis approach. *Science of the Total Environment*, 2018, 610-611: 831-844.
- [64] Yidana, S.M., et al. Evolutionary analysis of groundwater flow: Application of multivariate statistical analysis to hydrochemical data in the Densu Basin, Ghana. *Journal of African Earth Sciences*, 2018, 138: 167-176.
- [65] Adomako, D., et al. Hydrogeochemical Evolution and Groundwater Flow in the Densu River Basin, Ghana. *Journal of Water Resource and Protection*, 2011, 03(07): 548-561.
- [66] Devic, G., D. Djordjevic, S. Sakan. Natural and anthropogenic factors affecting the groundwater quality in Serbia. *Sci Total Environ*, 2014, 468-469: 933-42.
- [67] Tran, D.A., et al. Hydrogeochemical characteristics of a multi-layered coastal aquifer system in the Mekong Delta, Vietnam. *Environmental Geochemistry and Health*, 2020, 42(2): 661-680.
- [68] Waziri, M., V.O. Ogugbuaja. Interrelationship between physicochemical water pollution indicators: A case study of River Yobe-Nigeria. *American Journal of Scientific and Industrial Research*, 2010, 1(1): 76-80.
- [69] Raji, M.I.O., et al. Physicochemical Characteristics of Water Samples Collected from River Sokoto, Northwestern Nigeria. *Atmospheric and Climate Sciences*, 2015, 05(03): 194-199.
- [70] Awoyemi, V.O., C.A. Achudume, A.A. Okoya. The Physicochemical Quality of Groundwater in Relation to Surface Water Pollution in Majidun Area of Ikorodu, Lagos State, Nigeria. *American Journal of Water Resources*, 2014, 2(5): 126-133.
- [71] UNICEF. Unicef Hand Book on Water Quality: United Nations Children's Fund, New York, 2008: 1-191. Accessed from: [https://www.unicef.org/french/wash/files/WQ\\_Handbook\\_final\\_signed\\_16\\_April\\_2008.pdf](https://www.unicef.org/french/wash/files/WQ_Handbook_final_signed_16_April_2008.pdf)
- [72] Mahananda, M.R., M. B.P., N.R. Behera. Physico-chemical Analysis of Surface and groundwater of Bargarh District, Orissa, India. *IJRRAS*, 2010, 2(3): 284-295.
- [73] Ramesh, R.P., et al. Water quality assessment of Begnas and Rupa Lakes, Lesser Himalaya Pokhara, Nepal. *Journal of the Institute of Engineering*, 2019, 15(2): 113-122.
- [74] Hamdy, A. Control of Agricultural water pollution in Kandiah, A. (Ed) Water pollution. CHEAM, International Conference Valenzano, Italy, 1997: 101-119.
- [75] Sankhla, M.S. Water Contamination through Pesticide & Their Toxic Effect on Human Health. *International Journal for Research in Applied Science and Engineering Technology*, 2018, 6(1): 967-970.
- [76] SON. Nigerian Industrial Standard (NIS 554:2007); The Nigerian Standard for Drinking Water Quality"

- Standard Organization of Nigeria, 2007: 1-28. Accessed from:  
<https://africacheck.org/wp-content/uploads/2018/06/Nigerian-Standard-for-Drinking-Water-Quality-NIS-554-2015.pdf>
- [77] Anderson, H.R., W. Ogilbee. Aquifers in the Sokoto basin, northwestern Nigeria, with a description of the general hydrogeology of the region: Contributions to the hydrology of Africa and the Mediterranean Region. Geological Survey Water-Supply Paper 1757-L, 1973: 1-88.
- [78] Wali, S.U., A.M. Bakari. Assessment of groundwater variability over different geologic formations across Kebbi State, Nigeria. *Zaria Geographer*, 2016, 3(1): 155-167.
- [79] Wali, S.U., et al. Groundwater Hydrochemical Characterization in Urban Areas of Southwestern Sokoto Basin Nigeria. *SF Journal of Environmental and Earth Science*, 2018, 1(1): 1-17.
- [80] Wali, S.U., et al. An examination of rural groundwater quality in southwestern Zuru Local Government Area, Kebbi State Nigeria. *The Nigerian Geographical Journal*, 2018, 12(1): 122-142.
- [81] Haritash, A.K., et al. Suitability assessment of groundwater for drinking, irrigation and industrial use in some North Indian villages. *Environmental Monitoring Assessment*, 2008, 145(1-3): 397-406.
- [82] Krainov, S.R., A.P. Belousova, B.N. Ryzhenko. Formation of high-alkali groundwater in upper aquifers. *Water Resources and Industry*, 2001, 28(5): 491-501.
- [83] Missi, C., E.A. Atekwana. Physical, chemical and isotopic characteristics of groundwater and surface water in the Lake Chilwa Basin, Malawi. *Journal of African Earth Sciences*, 2020, 162: 103737.
- [84] Scheelbeek, P.F., et al. Drinking Water Sodium and Elevated Blood Pressure of Healthy Pregnant Women in Salinity-Affected Coastal Areas. *Hypertension*, 2016, 68(2): 464-70.
- [85] Selvakumar, S., N. Chandrasekar, G. Kumar. Hydrogeochemical characteristics and groundwater contamination in the rapid urban development areas of Coimbatore, India. *Water Resources and Industry*, 2017, 17: 26-33.
- [86] Smith, L.E.D., G. Siciliano. A comprehensive review of constraints to improved management of fertilizers in China and mitigation of diffuse water pollution from agriculture. *Agriculture, Ecosystems & Environment*, 2015, 209: 15-25.
- [87] Deutsch, B., P. Kahle, M. Voss. Assessing the source of nitrate pollution in water using stable N and O isotopes. *Agronomy for Sustainable Development*, 2007, 26(4): 263-267.
- [88] El-Zeiny, A., S. El-Kafrawy. Assessment of water pollution induced by human activities in Burullus Lake using Landsat 8 operational land imager and GIS. *The Egyptian Journal of Remote Sensing and Space Science*, 2017, 20: S49-S56.